

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/006421

International filing date: 23 February 2005 (23.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/547,591  
Filing date: 24 February 2004 (24.02.2004)

Date of receipt at the International Bureau: 07 April 2005 (07.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1302290

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*March 31, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/547,591**

**FILING DATE: February 24, 2004**

**RELATED PCT APPLICATION NUMBER: PCT/US05/06421**



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

**PROVISIONAL APPLICATION COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION

022404  
16569 U.S. PTO

Docket No. 17596PROV (BOT)	Type a plus sign (+) inside this box →	+
----------------------------------	--	---

**INVENTOR(S)/APPLICANT(S)**

LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Fernandez-Salas	Ester		1710 Rocky Road Fullerton, CA 92831 Citizen of Spain
Garay	Patton	E.	2021 N. Beverly Plaza #137 Long Beach, CA 90815 Citizen of USA
Aoki	Kei Roger		2 Ginger Lily Court Coto de Caza, CA 92679 Citizen of USA

22264 U.S. PTO  
60/547591  
022404

**TITLE OF THE INVENTION (280 characters max)**

**SCREENING METHODS AND COMPOSITIONS**

**CORRESPONDENCE ADDRESS**

ALLERGAN, INC.  
Intellectual Property Law Department  
T2-7H  
2525 Dupont Drive  
Irvine

CA	Zip:	92612	Country:	U.S.A.
----	------	-------	----------	--------

**ENCLOSED APPLICATION PARTS (check all that apply)**

[ x ] Specification Number of Pages 66 [ ] Small Entity Statement

[ x ] Drawing(s) Number of Sheets 16 [ ] Other (specify):

**METHOD OF PAYMENT (check one)**


<input type="checkbox"/> A check of money order is enclosed to cover the Provisional filing fee	Provisional Filing Fee Amount (\$)	<b>\$160.00</b>
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number: 01-0885		

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☐ No

☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE  Date: 8/21/04

TYPED or PRINTED NAME: CARLOS A. FISHER, ESQ. REGISTRATION NO. 36,510

☐ Additional inventors are being named on separately numbered sheets attached hereto

**PROVISIONAL APPLICATION FILING ONLY**

SEND COMPLETED FORMS TO: COMMISSIONER FOR PATENTS, MAIL STOP PROVISIONAL PATENT APPLICATION, P.O. BOX 1450, ALEXANDRIA, VA 22313-1450

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: )  
Fernandez-Salas et al )  
Serial No.: applied for ) I hereby certify that this correspondence is being deposited  
) with the United States Postal Service as Express Mail No.  
Filed: Herewith ) (EV193718029US) addressed to:  
) Mail Stop Provisional Patent Application, Commissioner for Patents,  
) P.O. Box 1450, Alexandria, VA 22313-1450 on:  
For: SCREENING METHODS AND ) Date of Deposit: 2/24/2004  
COMPOSITIONS )  
) By: BONNIE FERGUSON  
) Signature: Bonnie Ferguson  
Examiner: Unknown ) Date of Signature: 2/24/2004  
\_\_\_\_\_ )

TRANSMITTAL SHEET


Commissioner for Patents  
Alexandria, VA 22313-1450

Sir:

Enclosed herewith are the following documents:

- (1) Return/postage paid Postcard
- (2) Transmittal Sheet
- (3) Provisional application Cover Sheet (1 pages)
- (4) Specification ( 66 pages)
- (5) Drawings (16 sheets)

Respectfully Submitted,

  
Bonnie Ferguson

**SCREENING METHODS AND COMPOSITIONS**

Field of the Invention

5           The present invention is relevant to the fields of  
medicine, molecular biology, biochemistry and medicinal  
chemistry.

Background of the Invention

10

          Neurotoxins, such as those obtained from  
*Clostridium botulinum* and *Clostridium tetanus*, are  
highly potent and specific poisons of neural cells.  
These Gram positive bacteria secrete two related but  
15   distinct toxins, each comprising two disulfide-linked  
amino acid chains: a light chain (L) of about 50 KDa  
and a heavy chain (H) of about 100 KDa, which are  
wholly responsible for the symptoms of these diseases.  
Proteolytic cleavage of the H chain gives rise to two  
20   fragments of approximately 50 KDa each: a carboxy  
terminal domain (H<sub>C</sub>) which contains the cell surface-  
binding domain, and an amino terminal domain (H<sub>N</sub>) which  
contains a region responsible for translocation of the  
toxin.

25

          The tetanus and botulinum toxins are among the  
most lethal substances known to man, having a lethal  
dose in humans of between 0.1 ng and 1 ng per kilogram  
of body weight. Tonello et al., *Adv. Exp. Med. & Biol.*

389:251-260 (1996). Both toxins function by inhibiting

neurotransmitter release in affected neurons. The tetanus neurotoxin (TeNT) acts mainly in the central nervous system, while botulinum neurotoxin (BoNT) acts

5 at the neuromuscular junction by inhibiting acetylcholine release from the axon of the affected neuron into the synapse, resulting in a localized flaccid paralysis. The effect of intoxication on the affected neuron is long-lasting and has been thought to  
10 be irreversible.

The tetanus neurotoxin (TeNT) is known to exist in one immunologically distinct type; the botulinum neurotoxins (BoNT) are known to occur in seven  
15 different immunogenic types, termed BoNT/A through BoNT/G. While all of these types are produced by isolates of *C. botulinum*, two other species, *C. baratii* and *C. butyricum* also produce toxins similar to /F and /E, respectively. See e.g., Coffield et al., *The Site and Mechanism of Action of Botulinum Neurotoxin in Therapy with Botulinum Toxin* 3-13 (Jankovic J. & Hallett M. eds. 1994), the disclosure of which is  
20 incorporated herein by reference.

25 It is also recognized by those of skill in the art that within each type of clostridial neurotoxin there can be various strains varying somewhat in their amino acid sequence, and also in the nucleic acids encoding these proteins. Figures 16A-16C show an alignment of  
30 various known strains of BONT/A showing their amino

acid sequences and areas of difference. Additional BONT/A-producing strains are known in the art, such as, without limitation, strain NCTC 2012, and those listed in Franciosa et al., 32 J. Clin. Microbiol. 1911

5 (August 1994). BONT/A neurotoxin isolated from these strains may contain conservatively modified variations of their amino acid sequences and of the nucleotide sequences encoding them.

10           Regardless of type, the molecular mechanism of intoxication appears to be similar, although all details of this process are not yet precisely known and Applicants have no wish to be limited by the following description.

15           In the first step of the process, the toxin binds to the presynaptic membrane of the target neuron through a specific interaction between the heavy chain and a cell surface receptor; this receptor is thought  
20 to be different for each type of botulinum toxin and for TeNT. The carboxy terminus of the toxin heavy chain appears to be required for targeting of the toxin to the cell surface.

25           In the second step, the toxin crosses the plasma membrane of the poisoned cell. The toxin is first engulfed by the cell through receptor-mediated endocytosis, and an endosome containing the toxin is formed. The catalytic domain of the toxin is then  
30 translocated from the endosome into the cytoplasm of

the cell. This last step is thought to be mediated by the amino terminus of the heavy chain, which undergoes a conformational change in response to a pH of about 5.5 or lower. Acidic endosomes are known to possess a proton pump which decreases intra-endosomal pH. The conformational shift is believed to expose hydrophobic residues in the translocation domain, which permits the toxin to embed itself in the endosomal membrane. The catalytic domain then translocates through the endosomal membrane, probably via a pore, into the cytosol.

The last step of the mechanism of botulinum toxin activity appears to involve reduction of the disulfide bond joining the heavy and light chain. The entire toxic activity of botulinum and tetanus toxins is contained in the light chain of the holotoxin; the light chain is a zinc ( $\text{Zn}^{++}$ )-dependent endopeptidase.

In neurons, neurotransmitters are packaged within synaptic vesicles, formed within the cytoplasm, then transported to the inner plasma membrane where the vesicles dock and fuse with the plasma membrane. Recent studies of nerve cells employing clostridial neurotoxins as probes of membrane fusion have revealed that fusion of synaptic vesicles with the cell membrane in nerve cells depends upon the presence of specific proteins that are associated with either the vesicle or the target membrane. See *id.* These proteins have been termed SNAREs. A protein alternatively termed



synaptobrevin or VAMP (vesicle-associated membrane protein) is a vesicle-associated SNARE (v-SNARE).

There are at least two isoforms of synaptobrevin; these two isoforms are differentially expressed in the mammalian central nervous system, and are selectively associated with synaptic vesicles in neurons and secretory organelles in neuroendocrine cells. The target membrane-associated SNAREs (t-SNAREs) include syntaxin and SNAP-25. Following docking, the VAMP protein forms a core complex with syntaxin and SNAP-25; the formation of the core complex appears to be an essential step to membrane fusion. See Rizo & Sudhof, *id.* and Neimmann et al., *Trends in Cell Biol.* 4:179-185 (May 1994).

15

The endopeptidase of clostridial toxin light chain selectively cleaves SNARE proteins. TxNT, BoNT/B, BoNT/D, BoNT/F, and BoNT/G intoxication causes cleavage of VAMP. Most of the VAMP present at the cytosolic surface of the synaptic vesicle is removed as a result of any one of these cleavage events. Each toxin specifically cleaves a different bond and/or substrate, except TeNT and BoNT/B, which cleave the same bond in the VAMP protein.

25

BoNT/A and /E selectively cleave the plasma membrane-associated protein SNAP-25; this protein is bound to and present on the cytosolic surface of the plasma membrane. BoNT/E cleaves a 26-amino acid fragment from the C terminus of SNAP-25 whereas BoNT/A

30

removes only nine residues, creating a 197-amino acid  
fragment (SNAP25<sub>197</sub>) that is 95% the length of SNAP-25.  
BoNT/C cleaves syntaxin, an integral protein having  
most of its mass exposed to the cytosol. Syntaxin  
5 interacts with the calcium channels at presynaptic  
terminal active zones. See Tonello et al., *Tetanus and  
Botulism Neurotoxins in Intracellular Protein  
Catabolism* 251-260 (Suzuki K & Bond J. eds. 1996).  
BoNT/C<sub>1</sub> also cleaves SNAP-25 at a peptide bond next to  
10 that cleaved by BoNT/A.

Both TeNT and BoNT are taken up at the  
neuromuscular junction. BoNT remains within peripheral  
neurons, and blocks release of the neurotransmitter  
15 acetylcholine from these cells. Through its receptor,  
TeNT enters vesicles that move in a retrograde manner  
along the axon to the soma, and is discharged into the  
intersynaptic space between motor neurons and the  
inhibitory neurons of the spinal cord. At this point,  
20 TeNT binds receptors of the inhibitory neurons, is  
again internalized, and the light chain enters the  
cytosol to block the release of the inhibitory  
neurotransmitters 4-aminobutyric acid (GABA) and  
glycine from these cells. Id.

25

Because of its specifically localized effects,  
dilute preparations of BoNT have been used since 1981  
as therapeutic agents in the treatment of patients  
having various spastic conditions, including strabismus  
30 (misalignment of the eye), blepharospasm (involuntary

eyelid closure) and hemifacial spasm. See e.g.,

Borrodic et al., *Pharmacology and Histology Botulinum Toxin in Therapy with Botulinum Toxin* 3-13 (Jankovic J. & Hallett M. eds. 1994). Of the seven toxin types,

5 BoNT/A is the most potent of the BoNTs, and the most well characterized.

Intramuscular injection of spastic tissue with dilute preparations of BoNT/A has also been used  
10 effectively to treat spasticity due to brain injury, spinal cord injury, stroke, multiple sclerosis and cerebral palsy. The extent of paralysis depends on both the dose and dose volume delivered to the target site. Typically, the neurotoxin is administered in a  
15 preparation that also contains several non-toxic proteins as well, including hemagglutins and associated glycoproteins that assist in maximizing its stability and presentation to the target motor neuron.

20 Recently, it has been suggested that BONT's and TeNT bind neurons with through a high-affinity interaction between the toxin binding domain (Hc), a ganglioside, and an unknown protein receptor at the nerve terminal. See e.g., Schiavo et al., 80 *Physiol. Rev.* 717 (2000), Williamson, 274 *J. Biol. Chem.* 25173  
25 (1999), Chaddock, *Clostridium Botulinum And Associated Neurotoxins in Molecular Medical Microbiology* 1141 (Academic Press 2001). Although toxin binding is not strictly dependent on the presence of gangliosides, the  
30 presence of specific gangliosides appears to be

required for high affinity binding. In particular,  
BONTs have been observed to interact *in vitro* and *in vivo* with polysialogangliosides, especially those of  
the Glb series (GD1b, GT1b and Gq1b) Halpern & Neale,  
5 195 *Curr. Topics Microbiol.* 221 (1995).

A Japanese group and a research group from the  
University of Wisconsin have reported the results of  
experiments that suggest the identification of a neural  
10 cell membrane receptor for BONT/B and the involvement  
of ganglioside GT1b in binding of this neurotoxin to  
its target. See Kozaki S., 25 *Microb. Pathog.* 91-99  
(Aug. 1998); Dong, M. et al., *J. Cell Biol.*  
62:1293 (September 29, 2003). While these results  
15 appear to show that fragments of synaptotagmin I and  
synaptotagmin II bind BONT/B, the results are largely  
circumstantial. Indeed, there is later evidence which  
intimates that these proteins may simply be chaperone  
proteins for the neurotoxin. Coffield and  
20 Kalandakanond, Abstract 29 page R16.

Thus, to date the identity of the membrane  
receptor for any of the clostridial neurotoxins has not  
been conclusively demonstrated.

25

#### Summary of the Invention

The present invention is based on the  
identification of a neural cell membrane receptor to  
30 which the BONT/A neurotoxin selectively binds as the

first step to the selective intoxication of the neuron.

The present inventors have also identified specific gangliosides which facilitate binding of BONT/A to the FGFR3 receptor and the internalization of the BONT/A

5 and BONT/E neurotoxins within the neural cell.

All of the publications cited in this application are hereby incorporated by reference herein.

10 Brief Description of the Drawings

Figure 1 is a graph showing the decrease in insulin secretion from HIT-T15 cells given PURE-A toxin by electroporation. As can be seen, the addition of glucose to 25 mM induced insulin secretion from untreated cells and cells subjected to electroporation without the addition of PURE-A. However, cells into which PURE-A was introduced were unresponsive to  
15  
20 induction of insulin secretion.

Figure 2A demonstrates that the presence of the toxin delayed growth in HIT-T15 cells when compared to controls, but toxin-treated cells were able to replicate normally after a recovery period. Figures 2B and 2C demonstrate that cleavage of SNAP-25 was detected by Western blot at all time points tested when PURE-A was introduced into the cells.  
25

Figure 3 shows HIT-T15 cells, transformed with a human brain cDNA library and selected using magnetic beads to which BONT/A had been bound. Individual colonies are visible in the dish and are surrounded by magnetic beads.

Figure 4 shows the results of an assay of insulin release from HIT-T15 cells containing the putative BONT/A receptor. Cells were exposed to 1 nM PURE-A and inhibition of insulin release upon glucose stimulation was measured using an enzyme-linked immunoassay.

Figure 5 shows the reduction of insulin release in representative HIT-T15 transformants C6 and C7 upon incubation with BONT/A.

Figure 6 shows Western blots of clones C6 and C7 incubated with BONT/A using anti SNAP-25 antibody.

Figure 7A shows the results of a PCR amplification of a region bordering the human brain library cDNA insert in the c7 cell line.

Figure 7B shows a further amplification of from the C7 cDNA primer using a set of nested primers.

Figure 8 shows the levels of SNAP-25<sub>197</sub> produced upon incubation of three cell lines (Neuro-2A, SH-SY5Y, and NG108-15) with BONT/A.

Figure 9A shows the time course of the appearance of the SNAP<sub>197</sub> fragment in Western blots of Neuro 2 cell lysates following BONT/A treatment. Figure 9B shows  
5 the same experiment in SH-SY5Y cells.

Figure 10 shows the cooperative effect of ganglioside GT1b on the rate and/or extent of SNAP-25 cleavage in Neuro-2A cells treated with 12.5, 25 and 50 pM BONT/A.

10 Figure 11 the overall reaction scheme to crosslink BONT/A with the putative receptor in Neuro-2A cells using the cross-linking reagent sulfo-BED.

Figure 12A illustrates the isolation of a complex  
15 of approximately 250 kDa from Neuro-2A cells containing the 150 kDa neurotoxin cross-linked to the putative BONT/A receptor. Bands were visualized with silver staining.

20 Figure 12B shows the same sample, electrophoresed on a reducing gel to separate toxin from receptor, then transferred to a membrane for Western blotting. Subsequent probing of the membrane with hyperimmune neutralizing serum to PURE A results in two bands: one  
25 corresponding to the holotoxin (150 kDa) and the other to heavy chain (100 kDa). In this Figure, the stronger binding of the serum to the heavy chain in preference

to the light chain results in preferential detection of the heavy chain rather than the light chain.

Figures 13A-D shows a Western blot of the membrane-transferred proteins from the reducing gel with antibodies selective for each of FGFR1, FGFR2, FGFR3, and FGFR4, respectively. Only antibodies selective for FGFR3 (Figure 13C) bound to bands in the Western blot.

Figure 14 A-D shows Western blots taken from lysates from Neuro-2A, SH-SY5Y, PC-12, HIT-T15, and HIT-T15 clone C7 subjected to SDS-PAGE. The membrane-transferred proteins were evaluated for the presence of FGFR1-4 using antibodies selective for each of these FGF receptors (Figures 14 A-D, respectively).

Figure 15 shows the results of a Western blot of a receptor competition experiment in Neuro-2 cells of PURE-A and FGF.

Figure 16A-C shows an alignment of some representative strains of BONT/A.

## Detailed Description of the Invention

The present invention is based in part on the finding that high affinity binding of neurotoxins, particularly (though not exclusively) BONT/A, to neuronal cells requires a ganglioside component and a



protein receptor component. To this end, the present inventors have first discovered that the fibroblast growth factor 3 (FGF3) receptor, when, is sufficient to bind and direct the intracellular internalization of BONT/A. Secondly, the inventors have discovered that when the FGFR3 is combined with gangliosides, particularly the ganglioside GT1b, the affinity of both toxin binding and internalization is significantly increased.

10

The C-terminal region of the heavy chain of BONT/A (termed the Hc region), like all known clostridial neurotoxins, is involved in specific binding to a neuronal receptor present on the external surface of neurons.

15

The Hc binding domain is composed of two distinct structural domains (reviewed in Poli, M.A. and Lebeda F.J., *An overview of Clostridial Neurotoxins* in I Handbook of Neurotoxicology 293-304 (E.J. Massaro ed., Humana Press, 2002), and Meunier et al., *Molecular mechanism of action of botulinal neurotoxins and the synaptic remodeling they induce in vivo at the skeletal neuromuscular junction* in I Handbook of Neurotoxicology 305-347 (E.J. Massaro ed., Humana Press, 2002). The N-terminus of the Hc region presents a jelly-roll architecture related to that of the S-lectins, a carbohydrate-binding family of proteins. By contrast, the C-terminus of Hc is in a pseudo threefold trefoil conformation that presents structural similarity to the

30

sequentially unrelated interleukins- $1\alpha$  and  $1\beta$ , Kunitz-type trypsin inhibitors, as well as fibroblast growth factors (FGF). These proteins, mostly  $\beta$ -proteins, are involved in protein-protein interactions. This analysis is therefore consistent with the hypothesis that clostridial neurotoxins bind to receptors comprised of two components; a protein component and a carbohydrate component.

10       The amino acid sequence at the C-terminus of Hc is poorly conserved among different clostridial neurotoxins, and competition experiments have shown that different BoNT serotypes bind to different protein receptors on the surface of neuronal cells. Further, BoNTs are known to interact *in vivo* and *in vitro* with polysialogangliosides, in particular with members of the Glb series (GD1b, GT1b, and Gq1b) (reviewed in Halpern and Neale, 195 *Current Topics Microbiol. Immunol.*, 221-241 (1995)). Preincubation of the toxin with these gangliosides protects the neuromuscular junction (NMJ) of mice from BoNT toxicity.

High-affinity, trypsin-sensitive, BoNT-binding sites were found in isolated synaptosomes (Williams et al, 131 *Eur. J. Biochem.* 437-445 (1983)). Since lectins with high affinity for sialic acid antagonize the binding of BoNTs, their protein receptors may be glycoproteins. Receptors for BoNTs would direct them to acidic vesicles allowing the translocation of the LC into the cytosol of the neuron.

Based on these findings, and as the result of the data described herein, the Applicants have discovered that the fibroblast growth factor receptor 3 (FGFR3) binds BONT/A on the surface of neural cell lines.

5 Internalization of the toxin can be followed when these cell lines are exposed to the toxin. Moreover, BONT/A internalization is inhibited in a dose-dependent manner when FGF, preferably an FGF selected from FGF 1, 2, and 9, is added at increasing concentrations. Cells tested  
10 by the Applicants that did not display the FGFR3 receptor were unable to internalize the toxin, although when subjected to electroporation in the presence of BONT/A, the intracellular cleavage of SNAP-25 could be detected, indicating that the endopeptidase activity of  
15 the toxin remained intact, and that the cells remained susceptible to the endopeptidase.

Accordingly, in one embodiment the present invention is directed to a method of screening a sample  
20 for the presence of a molecule able to compete with BONT/A for selective binding to neural cells of the neural muscular junction comprising:

- a) contacting said sample with a composition comprising FGFR3 receptor and, optionally,  
25 GT1b ganglioside, and
  - b) determining whether said molecule selectively binds FGFR3,
- wherein selective binding of said molecule to FGFR3 indicates that said molecule is able to inhibit or  
30 reduce BONT/A toxicity to said neural cells.

By "BONT/A" or "type A" or similar terminology referring unambiguously to *Clostridium botulinum* neurotoxin type A is meant any of a number of  
5 polypeptide neurotoxins, and derivatives thereof, which can be purified from *Clostridium botulinum* serotype A strains and which share FGFR3 as a cell surface receptor. Such neurotoxins include those found in or corresponding to the following strains and accession  
10 numbers:

	<u>Strain</u>	<u>accession numbers</u>
	CL138	AAQ16535
15	137	AAQ16534
	129	AAQ16533
	13	AAQ16532
	42N	AAQ16531
	Hall A-hyper	AAM75961
20	667Ab	CAA61124
	NCTC 2916, P10845	CAA36289
	Allergan-Hall A	AAQ06331
	62A	AAA23262
	Kyoto-F	CAA51824
25	type A NIH	BAA11051
	NCTC 7272	
	7I03-H	
	Kumgo	AA021363

30 Additionally, BONT/A includes those polypeptides whose amino acid sequences are listed in Figure 16 hereof.

The molecule to be tested in the screening method may be a "small" organic compound of synthetic origin,  
35 or may be a macromolecule (either of synthetic or biological origin), such as a polypeptide (including

without limitation, a growth factor, a neurotoxin, a modified neurotoxin, an antibody or an antibody derivative), nucleic acid (such as a nucleic acid aptomer), or polysaccharide (such as a ganglioside or a lectin). In one preferred embodiment, the molecule is a synthetic molecule designed based on the tertiary structure and three dimensional conformation of FGF or an antibody that inhibits BONT/A binding to the FGF receptor. Such SAR (structure/activity relationship) analysis is routine in the art of medicinal chemistry, among other fields.

Determination of whether the molecule selectively binds the FGFR3 can be accomplished in any appropriate manner. For example, and without limitation, the molecule may be labeled with a detectable moiety, such as a radionuclide (e.g.,  $^{14}\text{C}$ ,  $^3\text{H}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$  or the like), a fluorescent moiety, or with an affinity tag, such as poly histadine, biotin, or GST (glutathione-S-transferase). The latter tags, which are common in the art and can be created as part of a fusion protein, can be used to purify the receptor/test compound complex.

By "selective" binding is mean that a binding agent is able to bind its target under physiological conditions, or in vitro conditions substantially approximating physiological conditions, to a statistically significantly greater degree (i.e., has a smaller  $K_d$  or dissociation constant) than to other, non-target molecules on the surface of the neural cell.

"K<sub>d</sub>" is the molar concentration of the binding agent at which half the target molecules are bound by the binding agent.

5           Alternatively, the binding of the molecule and receptor can be detected using labeled or unlabeled antibodies selective for (as an example) the molecule (test compound), FGFR3, or the ganglioside (such as GT1b). If the antibody is labeled, the binding of the  
10 molecule can be detected by various means, including Western blotting, direct microscopic observation of the cellular location of the antibody, measurement of cell or substrate-bound antibody following a wash step, or electrophoresis, employing techniques well-known to  
15 those of skill in the art. If the antibody is unlabeled, one may employ a labeled secondary antibody for indirect detection of the bound molecule, and detection can proceed as for a labeled antibody.

20           Other means of detecting the binding of the molecule include competition experiments, whereby the molecule is used to test for inhibition of FGF or BONT/A binding to the FGFR3, either directly or indirectly. Indirect means may include detection of  
25 the effect of the molecule (test compound) on the activity of FGF or BONT/A, for example, by detecting the cleavage of full-length SNAP-25 to SNAP-25<sub>197</sub> by the neurotoxin.

In another embodiment, the invention is drawn to a polypeptide comprising at least the H<sub>c</sub> region of BONT/A, which is produced from a bulk or formulated preparation wherein the bulk or formulated preparation is assayed for specific binding to neural cells using a method comprising contacting said polypeptide with a composition comprising FGFR3 receptor and, optionally, GT1b ganglioside, and determining whether said polypeptide selectively binds FGFR3.

10

In another embodiment similar to the above aspect of the invention, the polypeptide comprises at least an FGFR3 binding domain, other than the H<sub>c</sub> domain of BONT/A. Such a binding domain may comprise, for example, an FGF, such as FGF 1, FGF2, FGF4 or FGF 9, or an anti-FGFR3 antibody. Further, the polypeptide may optionally contain a translocation domain such as the H<sub>N</sub> domain of BONT/A. Additionally, the polypeptide will generally contain a clostridial neurotoxin light chain or variation thereof - the nature and/or source of the light chain can provide differences in the extent and half-life of the therapeutic effect of the polypeptide.

20

Thus, in this embodiment the claimed polypeptide is produced (which production may include purification, enzymatic treatment, and/or oxidation steps) from a bulk or formulation preparation. In one preferred embodiment the preparation may be, for example, a cell lysate from fermentation of a BONT/A-producing strain

25

30

of *Clostridium botulinum*, or from a suitable mammalian, insect or bacterial host cell producing a recombinant version of BONT/A. Such a bulk preparation may also be produced using cell-free transcription methodologies.

5 In another embodiment the preparation may be purified BONT/A formulated with associated stabilizing proteins, such as serum albumin. In each case, the preparation may comprise BONT/A molecules which are denatured or otherwise incorrectly folded so as not to bind to the  
10 target cells. The potency and/or specific activity of the preparation, or of fractions purified from the preparation, can be detected by using the claimed assay method.

15 Alternatively, the polypeptide to be assayed may comprise only a portion of the entire BONT/A molecule. For example, the bulk preparation may contain only the heavy chain of BONT/A, as separate production of the heavy and light chains of the toxin may be a preferred  
20 way of avoiding accidental exposure to the neurotoxin by laboratory workers.

As another example of the above embodiment, the polypeptide may comprise a chimeric recombinant  
25 polypeptide which contains the Hc region of the heavy chain of BONT/A (or some other FGFR3-binding moiety, such as FGF itself). The chimeric polypeptide comprises amino acid sequence regions additional to, or other than, those present in the wild-type BONT/A  
30 BONT/A molecule. For example, as disclosed in US



Patent No. 6,203,794, hereby incorporated by reference herein, botulinum and tetanus toxins may be used as the basis for the creation of transport proteins. The light chain of these transport proteins are generally  
5 either replaced by a therapeutic moiety or inactivated and coupled to such a therapeutic moiety. Additionally, International Patent Application WO0114570, published March 1, 2001, discloses chimeric neurotoxins comprising polypeptides containing domains of more than  
10 one neurotoxin. Thus, this aspect of the invention also encompasses, as a preferred embodiment, chimeric neurotoxins containing at least the Hc domain of BONT/A. Such molecules may be useful in modulating the time or extent of the inhibition of secretory vesicle  
15 release. Further, it may be desirable to target agents, such as therapeutic agents, to the extracellular surface of the neural cell membrane. Thus, such an agent may be joined (e.g., as a fusion protein or via post translational conjugation) to the  
20 Hc portion of BONT/A. In such a case the cell lysate or conjugation reaction mixture may comprise a batch preparation in accordance with this aspect of the invention.

25       The above-referenced polypeptides are screened for binding and/or internalization essentially as mentioned above in the described screening method embodiment.

      In yet another embodiment, the present invention  
30 is drawn to a method of marketing a polypeptide which

contains a region capable of binding the FGFR3 receptor comprising:

- 5           a)           obtaining permission from a governmental  
                  or regional drug regulatory authority to  
                  sell said polypeptide, wherein said  
                  polypeptide is first produced from a  
                  bulk preparation which is assayed for  
                  selective binding of said polypeptide to  
10           neural cells by contacting the bulk  
                  preparation containing said polypeptide  
                  with a composition comprising FGFR3  
                  receptor, and optionally GT1b  
                  ganglioside, and determining whether  
15           said polypeptide selectively binds FGFR3  
                  under such conditions,
- b)           packaging said polypeptide for sale in a  
                  manner consistent with the requirements  
                  of said regulatory authority, and
- 20           c)           offering said polypeptide for sale.

             In this embodiment the invention is drawn to a  
method of marketing a polypeptide containing the H<sub>c</sub>  
region of a BONT/A toxin. The polypeptide at issue in  
25   this embodiment of the invention is produced from a  
bulk preparation which is assayed for purity or  
activity using the screening method described  
previously. In a step of this method, permission is  
obtained from a regulatory body for the marketing of  
30   such polypeptide. In this context "permission" may be

tacit or express; that is, permission or approval may be obtained from the regulatory authority for the sale of a therapeutic agent or composition comprising said polypeptide, in which case "permission" is marketing  
5 approval for the sale of such agent or composition. Alternatively, "permission", as used herein, may comprise the assent, either affirmatively given or manifested by its lack of objection, of such regulatory authority to the continued sale of a product containing  
10 a polypeptide assayed in this new manner. As before, the polypeptide may comprise BONT/A, or a derivative thereof, or a fusion protein or conjugate containing the H<sub>c</sub> region of the BONT/A heavy chain.

15 By a "sample" or "preparation" is meant a preparation containing a agent or compound to be tested for its ability to bind to the BONT/A receptor. As used herein, a sample includes, without limitation, a solution comprising a small molecule, an antibody  
20 preparation, a cell lysate, a cell culture medium, a bulk or formulation preparation of a therapeutic agent, or a fraction from a purification process.

By an "agent" able to compete with BONT/A for  
25 selective cell binding is meant a compound, molecule, multimer, complex, or other composition which has the same cell surface target as BONT/A.

By "LD<sub>50</sub> assay" is meant an *in vivo* assay of neurotoxin activity comprising determining the dose of neurotoxin at which 50% of treated animals die.

5 Unless the context clearly indicates otherwise, by a "polypeptide", is meant a composition comprising one or more associated polymeric chains of amino acids linked by peptide bonds. If comprised more than one chain, the chains may be linked or otherwise associated  
10 by interactions such as disulfide bonds or ionic or hydrophobic interactions.

The therapeutic product comprising the polypeptide originally contained in the bulk preparation so assayed  
15 is labeled in accordance with the requirements of the regulatory authority. The product is then offered for sale. Offering for sale may comprise advertising or sales activity, educational seminars directed at doctors, hospitals, insurers, or patients,  
20 conversations with state, regional or governmental officials concerning subsidy reimbursement (such as Medicare or Medical).

## 25 Identification of BONT/A Receptor and Examples

We utilized two approaches for the identification of BoNT/A receptor: a) genetic complementation, and b) isolation from cells able to take up the BONT/A toxin  
30 with high affinity.

Cloning by genetic complementation is achieved by transfecting cells that do not contain the receptor with a vector containing a retroviral cDNA library  
5 constructed from a tissue known to express the receptor. Selection of cells expressing the receptor from a pool of transfected cells can be achieved by several means, such as binding the selected protein, or measuring its biological effect. According to this  
10 strategy, the receptor is then identified by PCR cloning of cDNA obtained from cells expressing the receptor.

The second approach, isolation of the putative  
15 receptor from cells known to expressing the receptor, required the identification of one or more cell line expressing a high affinity BONT/A receptor. Once identified the bound toxin is cross-linked with the receptor *in situ* and the cross-linked complex is  
20 isolated. Identification of the putative receptor can then be achieved by any number of means, including (without limitation) liquid chromatography-mass spectroscopy (LC-MS), or by Western blots using antibodies selected based on information on the  
25 literature which might offer clues as to the identity of the putative receptor.

Using both approaches we have identified FGFR3 as a receptor system enabling BoNT/A to selectively bind  
30 and enter neurons and other BONT/A susceptible cells.

Since the conformation of the binding domain of the various BoNT serotypes are structurally very similar (although the precise amino acid sequences vary) we also believe that serotypes other than BoNT/A may be able to use other members of the FGFR family (or other closely related members of the tyrosine kinase receptor superfamily, such as, without limitation, FGFR1, 2, 4 or 5 the PDGF receptor, and the EGF and VEGF receptors (including any isoforms) as a receptor in the plasma membrane of neuronal and other susceptible cells. Such other cells include chromaffin cells, acinar cells, and insulinoma cells. See Lomneth, 18 *J. Toxicol. Toxin Rev.* 77-94 (1999).

**I. Cloning of BoNT/A receptor by Genetic Complementation**

We first undertook the cloning of the neurotoxin receptor by genetic complementation. This procedure involves transferring the DNA for the receptor to a cell line that does not contain the receptor naturally. For this latter purpose we choose the insulinoma HIT-T15 cells. HIT-T15 cells secrete insulin in response to high levels of glucose in the medium, however BONT is unable to bind to these cells; thus extracellular exposure of HIT-T15 cells to BONT does not effect the secretion of glucose.

Despite the fact that HIT-T15 cells are normally unaffected by BONT, the intracellular secretory

exocytic process of these cells depends on the presence of SNAREs (which are targets of the neurotoxin endopeptidase activity) for vesicle docking and fusion. Specifically, HIT-T15 cells contain SNAP25, and  
5 intracellular introduction of BoNT/A by means not dependent upon toxin binding (such as permeabilization or electroporation) results in cleavage of the 206 amino acid full length SNAP25 molecule (SNAP25<sub>206</sub> into SNAP25<sub>197</sub>, the BONT/A specific cleavage product,  
10 resulting in failure of vesicle docking at the intracellular membrane surface, and the inhibition of insulin release.

Since neurons of the central nervous system are  
15 known to bind and internalize BONT/A, we used a human brain retroviral DNA library (purchased from Clontech, Inc.) as a source of the BONT/A receptor cDNA.

a) Receptor Cloning in HIT-T15 Cells

20

We packed the library into viruses using the Amphopack® 293 producer cell line. The retroviral library was then used to infect HIT-T15 cells. Cells assumed to be expressing the receptor clones were  
25 selected based on their ability to bind Dynex magnetic beads coated first with an antibody against the light chain of BONT/A, and then with pure BONT/A (PURE A); in this way the neurotoxin molecules were correctly oriented with the Hc (receptor binding domain) domain  
30 pointed outward.

Cells binding to the magnetic beads were isolated using a magnet. The cells were then plated in culture media and colonies permitted to develop in the dishes.

5 Individual colonies were picked and expanded to a 24 well plate to establish cell cultures. The isolated cells were grown and exposed to 1nM Pure BONT/A for 24 hours. Putative cell lines containing the receptor were selected based on the detection of SNAP-25<sub>197</sub> in

10 Western blot analysis (using anti-SNAP25<sub>197</sub> antibody) of whole cell lysates, and by the detection of the inhibition of insulin release.

Genomic DNA from cells testing positive for

15 cleavage of SNAP25 or inhibition of insulin release was isolated and used in PCR reactions using the Clontech ADVANTAGE® Genomic PCR kit and the following primers (written from 5' to 3'):

20 AGCCCTCACTCCTTCTCTAG  
SEQ ID No: 1

ACCTACAGGTGGGGTCTTTCATTCCC  
SEQ ID NO: 2

25 to amplify and isolate the human receptor gene present in the cells. Conditions were as recommended using the Clontech ADVANTAGE® Genomic PCR kit. Reactions were incubated at 95 degrees Celsius for 1 minute, followed

30 by 25 cycles at 68 degrees C for 30 seconds and 95



degrees C for 30 seconds, followed by 1 cycle at 68 degrees C for 6 minutes and final incubation at 4 degrees C.

5           A 12 kb fragment of DNA was identified and purified from the PCR reaction and subjected to a second PCR amplification. The primers used in the second PCR were nested primers designed to anneal to sequences of DNA within the amplicon originally  
10   produced, and had the following nucleotide sequences.

CCCTGGGTCAAGCCCTTTGTACACC

SEQ ID NO: 3

15

TGCCAAACCTACAGGTGGGGTCTTT

SEQ ID NO: 4

b) Inhibition of Insulin Secretion in HIT-T15 cells

20

The effect of BONT/A treatment on exocytosis was analyzed in HIT-T15 insulinoma cell transfectants selected as indicated above, and suspected to express the BONT/A receptor. These cells were treated for 15  
25   hours with 1nM pure BONT/A. Insulin secretion was measured in DMEM containing high glucose (25mM) or low glucose (5.6 mM). After 1 hour incubation with high glucose media at 37°C, the amount of insulin in the media was determined using an Insulin ELISA kit from  
30   Peninsula Laboratories. Exocytosis is expressed as the amount of insulin secreted per  $1 \times 10^5$  cells per hour.

II) Isolation Of The Putative Receptor From Cells Known  
To Express The BONT/A Receptor

5

a) Immunostaining PC12, HIT-T15, and Neuro-2 cells  
for Ganglioside GT-1b

10           Immunocytochemistry was performed on PC-12 (rat  
pheochromocytoma cells), Neuro-2 (mouse  
neuroblastoma), and HIT-T15 cell lines. These cells  
were fixed and probed with primary antibodies specific  
to the ganglioside GT-1b (obtained from Seikagaku  
15 Corp.) that is thought to be important for toxin  
uptake. We used Alexa<sup>®</sup>-fluor 488 (Molecular Probes,  
Inc.), conjugated secondary antibodies to visualize the  
GT-Ib via fluorescent microscopy. Those of skill in  
the art will recognize that Alexa<sup>®</sup>-fluor may be  
20 substituted with any fluorophore of similar intensity.

b) Identification of cell lines with high affinity  
uptake for BoNT/A

25           A method for neurotoxin treatment followed by  
Western Blot was developed to screen cell lines for  
toxin uptake using an antibody to detect the BONT/A  
cleavage product SNAP25<sub>197</sub>. Tested cell lines (human  
unless indicated otherwise) were: Neuro 2A, SH-SY<sub>5</sub>Y,  
30 NG108-15, N1E-115, SK-N-BE(2), SK-N-DZ, SK-N-F1, BE(2)-  
C, SK-N-SH, NB4 1A3, BE(2)-M17, C1300, NG108-15, HCN-  
1A, HCN-2 TE 189.T, ND8/34.

Cells of each candidate cell line were plated in poly-D-lysine/Laminin coated 6-well plates for 24 hours.

BoNT/A (900 KDa complex) was added to the cells at different concentrations (as described in the figures

5 and the results section below) in the culture medium overnight. Cells were then collected in 15 ml Falcon tubes, washed once with 1ml of PBS, and then transferred to 1.5 ml microcentrifuge tubes. Cells were lysed in 0.5 ml of lysis buffer (50 mM HEPES, 150 mM  
10 NaCl, 1.5 mM MgCl<sub>2</sub>, 1mM EGTA, 10% glycerol and 1% Triton X-100®) on a rotator at 4°C for 1 hour. Lysed cells were centrifuged at 5000 rpm at 4°C for 10 minutes to eliminate debris; supernatants were transferred to fresh siliconized tubes.

15

Protein concentrations were measured by Bradford's method and resuspended in 1 x SDS sample buffer at 1mg/ml or higher concentration. Samples were boiled for 5 min, 40 µl of the cell-free lysates were loaded on 4-  
20 12% Tris-HCl polyacrylamide gradient gels. The separated proteins were transferred to PVDF membranes for Western analysis and the membranes blocked in 5% non-fat milk in TBST buffer for 1 hour at room temperature. The cleaved SNAP25<sub>197</sub> was detected with  
25 antiSNAP25<sub>197</sub> antibody diluted in blocking buffer; blot was washed extensively, and the bound antibody was detected with a horseradish peroxidase-conjugated species-specific secondary antibody.

The Typhoon 9410 Imager (Amersham) for Western Blot Analysis was used instead of traditional film. After the final washes the membrane was reacted with ECL Plus Western blot detection reagent (Amersham), membrane was then incubated at room temperature for 5 min in order for the images to develop. The choice of pixel size and PMT voltage settings depended on each individual blot. Membranes were scanned and quantified using Typhoon Scanner and Imager Analysis software.

10

c) GT1b Ganglioside Treatment

Cells were cultured in Poly-D-lysine/Laminin coated 6-well plates in culture medium with 10% FCS for 24 hours. GT1b ganglioside (Alexis, San Diego, CA) was dissolved in PBS at 5mg/ml as a stock solution and stored at -20°C. At the time of treatment, GT1b was diluted in serum free medium to a final concentration of 25µg/ml and added to the cells in culture. The cells were incubated at 37°C, under 7.5% CO<sub>2</sub> for 24 hours. Following incubation, the GT1b-containing medium was removed and 0.5 ml to 5 ml BoNT/A complex was added at various concentrations (ranging from 0.125 nM to 5 nM) in culture medium (as recommended by the ATCC or ECACC tissue collections for that cell line) containing 10% FBS and incubated overnight.


20

25

d) Purification of the BoNT/A receptor from Neuro-2A Cells

30

Neuro-2 cells were found to have the most rapid toxin uptake profile (about 10 minutes) and affinity to the toxin, and were thus chosen for further study based on the presence of a high affinity toxin uptake system. Neuro-2 cells were therefore exposed to 5nM PURE-A at 4°C for 2-4 hours and subsequently treated with one of the two cross-linking agents BS<sup>3</sup> and sulfo-SBED (Pierce). The reagent BS<sup>3</sup> is an 11.4-Å chain water-soluble and membrane-impermeable agent that is not cleaved by reducing agents. The reagent sulfo-SBED contains three reactive groups (one of them designed to be UV-activated) and is designed to biotinylate a target protein. Its structure (including the presence of a biotin molecule) is given below:

**STRUCTURE OF SULFO-SBED**

The procedure for cross-linking the receptor and PURE-A was as follows.

5           All reactions are performed at 4 degrees C. 100  
ug of Pure A (0.34mL) is pelleted at 10,000 x g for 10  
minutes, and brought up in a final volume of 900 ul PBS  
(pH 7.4). The reaction is then transferred to the  
dark. Ten milligrams of Sulfo-SBED reagent is  
10 dissolved in a volume of 454 ul DMSO, to make a  
solution of 25 mM SBED. This is immediately serially  
diluted as follows: 100 ul of the 25 mM SBED reagent is  
added to 0.9 mL PBS (2.5 mM SBED)/ 10% DMSO). This in  
turn is diluted in the same manner to yield a 0.25 mM  
15 SBED)/ 1 % DMSO solution.

One hundred microliters of this latter solution is  
added to 900ul PURE A in PBS to yield 0.67 uM PURE A,  
25 uM Sulfo-SBED, 0.1% DMSO in 1 ml, and incubated in a  
20 4°C incubator for two hours in a secondary container on  
shaking apparatus. The reaction is stopped by the  
addition of 50 ul 1M TRIS pH 7.4 (final concentration  
50mM). The solution is inverted 6 times and allowed to  
incubate 30 minutes on ice. Meanwhile, Neuro-2A cells  
25 are harvested by centrifugation and washed 3 X with  
cold TBS, then divided into two aliquots of  $4 \times 10^8$   
cells. Each aliquot of cells is suspended in 12 mL  
cold TBS, and placed on ice for 15 minutes ( $3 \times 10^7$   
cells/mL). One milliliter of the Sulfo-SBED/PURE A  
30 solution is added to one of the Neuro-2A cell

suspension aliquots. The other aliquot receives the sulfo-SBED only (as a control). The final dilution thus contains  $4 \times 10^8$  cells ( $2 \times 10^7$  cells/mL) and 100 ug PURE A (33nM) in a 20 mL volume.

5

The PURE A/SBED and neuro-2 cell suspension is incubated at 4°C in an incubator for two hours in a secondary container on shaking apparatus. Each cell solution is distributed in 13 aliquots of 1.0 mL.

10 These aliquots are exposed to ultraviolet radiation (365 nm) for 15 minutes. The cells are harvested by centrifugation, washed with cold TBS, and lysed overnight at 4°C with 0.4 ml TRITON X-100 lysis buffer (50mM Hepes, 150mM NaCl, 1.5mM MgCl<sub>2</sub>, 1mM EGTA, 10%  
15 glycerol, 1% TRITON X-100, and suitable amounts of a protease inhibitor).

Following lysis, each aliquot is centrifuged at 5000 x g to remove unbroken cells, then the lysates  
20 each given 0.05mL of avidin-beads and incubated for 3 hours at 4°C. The avidin beads are then washed twice with 0.5mL lysis buffer (centrifuge at 1000 x g to pellet beads) followed by two washed with 0.5mL PBS buffer. The supernatant is then removed and combined  
25 with 100 ul SDS-PAGE loading buffer, then subjected to SDS-polyacrylamide gel electrophoresis.

e) Anti-FGFR antibodies used in Western blots

Antibodies to the cytoplasmic region of the polypeptides FGF 1 receptor (FGFR1), FGF 2 receptor (FGFR2), FGF 3 receptor (FGFR3) and FGF 4 receptor (FGFR4) were purchased from Santa Cruz Biotechnologies. The precipitated receptor-PureA complex (obtained as described above) was electrophoresed in an SDS polyacrylamide gel under reducing and non-reducing conditions. The resulting proteins were Western blotted and probed with the antibodies to FGFR1-FGFR4. Western blots were also prepared of cell lysates from Neuro-2A, HIT-T15 cells, HIT-T15C7 (i.e., the selected clone constructed as mentioned above and containing the putative receptor), SH-SY5Y, and PC12 cells to study the expression of the FGFR in the cell lines.

**f) FGF-Pure A Competition Experiments**

Neuro-2A cells were plated at a density of  $5 \times 10^5$  cells/well in 6 well plates the day before the experiment. PURE-A was added to the culture media at a final concentration of 5nM, either alone or in combination with FGF1, FGF2 or FGF1+FGF2 at various concentrations ranging from 0.1nM (physiological concentration for FGF) to 200nM. The treated cells were then incubated at 37°C for 10 min with the mixture. Cells were thereafter lysed and subjected to Western blot using an antibody to SNAP25 (SMI-81, from Sternburger Monoclonals) that recognizes both uncleaved and cleaved SNAP25.

**RESULTS**

**Strategy 1: Genetic Complementation**



The insulinoma cell line HIT-T15 secretes insulin upon glucose stimulation, depending on the presence of intact SNAP25. We decided to use this cell line for the genetic complementation because it permits the use of two separate indications of neurotoxin activity: Western blotting to detect the cleaved SNAP-25 (SNAP25<sub>197</sub>), and the detection of insulin secretion by monitoring insulin in the extracellular medium.

PURE-A was introduced into HIT-T15 cells by electroporation using a BioRad GENE PULSER® II to assess the cells' response by both parameters listed above before the receptor cloning started. Figure 1 shows that the presence of toxin inhibited exocytosis of insulin from cells induced to secrete insulin by the addition of 25 mM glucose.

Figure 2A shows the growth of HIT-T15 cells which were electroporated without and with PURE-A and their growth monitored for 10 days. Electroporation in the presence of PURE-A caused reduction in the cell number few days after cells were treated, but following this the growth of the cells was comparable to the control cells. Presence of toxin in the cells was followed by antibody detection of SNAP25<sub>197</sub> in the samples (C: control, E: electroporated without toxin, PA: electroporated with PURE-A). The antibody to SNAP25 in

Figure 2B detects cleaved and uncleaved products, while the antibody used in Figure 2C detects only SNAP25<sub>197</sub>.

To evaluate if the presence of the toxin in the cells affect cell growth, HIT-T15 cells were electroporated with PURE-A, and growth and presence of SNAP25<sub>197</sub> were monitored for 10 days. Figure 2A demonstrates that the presence of the toxin delayed growth when compared to controls, but toxin-treated cells were able to replicate normally after a recovery period. Figures 2B and 2C demonstrate that cleavage of SNAP25 was detected by Western blot at all time points tested when PURE-A was introduced into the cells.

Having determined that HIT-T15 cells were an appropriate cloning host for cloning the BONT/A receptor, these cells were transfected with media from AmphoPack<sup>®</sup> 293 producer cells containing a retroviral particles. The retrovirus used to infect the HIT-T15 cells contained the pRetroLIB<sup>®</sup> vector from Clontech. The cDNA human brain library inserts were cloned at the stuffer fragment using restriction endonuclease Sfi sites. The entire insert of these constructs, from the 5'LTR to the 3'LTR, incorporates into the genomic DNA of the host HIT-T15 cell.

Cells were cultured for four extra days to allow expression of the cDNA packaged in the virus vector. Selection of individual cells expressing putative

BONT/A receptors was achieved by incubating the cells for 30 min at 4°C with PURE-A attached to magnetic beads (Dynabeads®, Dynal Biotech). Cells able to bind PURE-A were selected with a magnet, washed, and plated again in dishes. Individual colonies were formed in the dish, as shown in Figure 3, which were surrounded by magnetic beads.

Those colonies were picked and cultured in 96 well plates. When individual wells were close to confluence, cells were exposed to 1 nM PURE-A and inhibition of insulin release upon glucose stimulation was measured using an enzyme-linked immunoassay (Fig. 4). As shown in Figure 4, HIT-T15 cells at later passage numbers have a reduced production and secretion of insulin; this is the reason why in the figure shows that the isolated clones' level of insulin release is lower than in the control cells, which were used at a lower passage.

20

Representative clones C6 and C7 demonstrated a reduction in the amount of insulin release in high glucose medium (25 mM) after treatment with 1 nM PURE-A (Fig. 5). The percentage of release is calculated using the amount of insulin released by cells exposed to toxin divided by the amount of insulin released by cells which were not exposed to toxin.

Expanded cultures of clones C6 and C7 were also exposed to PURE-A; cells were lysed and Western blots

performed to verify toxin uptake by immunological  
detection of SNAP25<sub>197</sub> (Figure 6).

The genomic DNA of the C7 cell line was used as a  
5 template in PCR reactions. The PCR primers used were  
LIB 5'® and LIB 3'® from Clontech - these have the  
nucleotide sequences labeled SEQ ID NO: 1 and 2 above,  
respectively; these primers anneal to vector sequences  
just upstream and downstream, respectively, of the  
10 cloned insert region. The PCR reaction resulted in the  
amplification of a 12-15 kb amplicon (Figure 7A, lane  
2).

We next performed a nested PCR reaction in an attempt to obtain smaller  
DNA fragments, as shown in Fig. 7B. Fragments were purified and cloned in  
15 Topo®-XL vectors (Invitrogen, Inc.). Plasmids containing inserts were partially  
sequenced, and analysis of the results demonstrated that the insert was derived  
from the retroviral vector.

**The neuroblastoma cell line Neuro-2A possesses the high affinity  
uptake system for BoNT/A**

20 We screened neuronal cell lines, (cell lines Neuro 2A, SH-SY5Y, N1E-  
115, SK-N-BE(2), SK-N-DZ, SK-N-F1, BE(2)-C, SK-N-SH, NB4 1A3, BE(2)-  
M17, C1300, NG108-15, HCN-1A, HCN-2 TE 189.T, and ND8/34) as described  
above, for their ability to uptake BoNT/A in cell culture. For the first step of the  
screening, cells were exposed to 1nM BoNT/A (900 kDa) overnight, and Western  
25 blots using antibodies selective for the presence of SNAP25<sub>197</sub> were performed.


Three cells lines: Neuro-2A, SH-SY5Y, and NG108-15  
displayed the highest levels of SNAP-25<sub>197</sub> following

BONT/A exposure (Fig.8) and were chosen for further evaluation.

The presence of the high affinity receptor is  
5 characterized by two attributes: rapid toxin uptake  
upon cell exposure, and the low toxin concentrations  
needed for intoxication. We evaluated the three  
selected cell lines for those criteria and discovered  
that Neuro-2A cells were able to take up toxin in less  
10 than 10 minutes, since a band for SNAP25<sub>197</sub> was seen ten  
minutes after treatment (Fig. 9A), the earliest time  
point tested. In contrast, the earliest point we  
detected the presence of SNAP-25<sub>197</sub> in SH-SY5Y cells was  
6 hours (Fig. 9B).

15

Toxin uptake was also increased by pretreating the  
cells with the complex ganglioside GT1b. This naturally  
occurring sialoglycolipid, which can be purchased from  
a number of sources, including Alexis Biosciences  
20 Corporation, Ltd.) has the following structure:



Neuro-2A cells pretreated with ganglioside can be  
affected by toxin at concentrations as low as 12.5pM  
5 (Fig. 10). The ganglioside GT1b has been described as  
the sugar moiety to bind to the neurotoxin binding  
domain in conjunction with binding to the unknown  
protein receptor. These data led us to conclude that  
Neuro-2A cells contain the high affinity uptake system  
10 for toxin, and were subsequently used to purify the  
receptor for BoNT/A.

Strategy 2: Isolation of the BoNT/A receptor from  
Neuro-2A cells

Neuro-2A cells were used as the source for the  
5 isolation of the protein receptor which binds BONT/A,  
as this cell line binds the toxin with the highest  
affinity among the cell lines tested, and internalizes  
the toxin rapidly. For this work we used the  
trifunctional sulfo-SBED reagent described above as a  
10 cross linking reagent. This reagent, after first being  
linked to biotin, will biotinylate a target protein  
receptor, which can then be subsequently precipitated  
using avidin as the affinity ligand.

15 As shown in Figure 11, the reagent is first linked  
to PURE-A through reaction with an amine-containing  
group in the toxin molecule. Then cells were exposed to  
5 nM PURE-A-sulfo-SBED complex at 4°C for 2-4 hours to  
avoid internalization of the toxin bound to receptor.  
20 Following binding of the receptor to the toxin, UV  
light was used to cross-link the neurotoxin and the  
receptor, and cells lysed.

The complex is isolated from the cell-free lysate  
25 using avidin beads and run on PAGE-SDS. Figure 12A  
shows the precipitated Neuro-2A protein visualized via  
silver stain. The SDS-PAGE was non-reduced, leaving  
the intact cross-linking reagent to bind PURE-A to  
putative receptor. The results show a large molecular  
30 protein complex (~250 kDa) that was isolated from cells

treated with BoNT/A, which would indicate the receptor has a size of approximately 100 kDa.

Figure 12B shows that the same sample was run on a reducing polyacrylamide gel to separate the toxin from the putative receptor. Following transfer of the reduced proteins to a membrane, the membrane was probed with antibodies to PURE-A. The resulting Western blot has two immunoreactive bands, which have molecular weights corresponding to the toxin (150 kDa) and heavy chain (100 kDa).

The membrane-transferred proteins from the reducing gel were also probed with antibodies selective for each of FGFR1, FGFR2, FGFR3, FGFR4, FGFR5 in an effort to more specifically identify the receptor to which the PURE A toxin bound. Figures 13A-D (corresponding respectively to FGF1, FGF2, FGF3, and FGF4) shows the resulting Western blots from these experiments. Only antibodies selective for FGF3 (Figure 13C) bound to bands in the Western blot. The main band has a molecular weight of approximately 80 KDa. However, higher molecular weight bands could also be detected, implying that the major band may be derived from a larger polypeptide.

We next surveyed a series of cell lines for the presence of the receptors FGFR1-4 in cell lysates. When indicated below, the presence of ganglioside GT1b was evaluated using an antibody to GT1b in fixed but non-



permeabilized cells. The cells used were as follow: a) Neuro-2A cells, which contain a high affinity BONT/A receptor protein and ganglioside GT1b, b) SH-SY5Y cells which contain a low affinity BONT/A receptor, c) 5 PC-12 cells, which contain a low affinity BONT/A receptor protein, and no ganglioside GT1b, d) HIT-T15 cells which contain ganglioside GT1b but no BONT/A receptor, e) HIT-T15 "C7" cells (obtained as explained above from exposure to retrovirus cDNA human brain 10 library), which are able to take up toxin (see Figure 5 (inhibition of insulin release) and Figure 6 (cleavage of SNAP 25)), and thus implicitly contain a BONT/A receptor.

15 Cell-free lysates were made from each of these cell lines (Neuro-2A, SH-SY5Y, PC-12, HIT-T15, and HIT-T15 clone C7) and were evaluated for the presence of the FGF receptors FGFR1, FGFR2, FGFR3, and FGFR4 through Western blots. (See Figures 14A-D, 20 respectively.)

The data from the Western blot revealed that FGFR2 and FGFR4 were not present in any of the cell lines. FGFR1 was present in all cell lines tested, including 25 the ones that do not take BONT/A, such as HIT-T15 (wt). FGFR3 was not present in the HIT-T15 wt cell line, but in contrast it was clearly present in the HIT-T15 C7 clone, generated from retroviral infection. This clone, upon expression of the cDNA inserted by the

retrovirus, is able to take up BoNT/A. The banding in HIT-T15 C7 was almost identical to that of Neuro-2A cells, but was different from the bands observed in SH-SY5Y cells that display low affinity BONT/A uptake.

5

To corroborate that BONT/A toxin enters Neuro-2A cells through the FGFR3 we performed a competition experiment with PURE-A at 5 nM concentration. Without added FGF, the toxin produces an almost complete

10 cleavage of SNAP25<sub>206</sub> into SNAP25<sub>197</sub> at this concentration. We added FGF1 and FGF2 at different concentrations ranging from 0.1 nM (the FGF concentration that produces its characteristic

15 biological effect) to 200 nM in order to be certain to obtain full competition. No ganglioside GT1b (other than that endogenous to the cell membranes of Neuro-2 cells) was added. The cells were exposed to either PURE A or PURE A plus FGF for 10 minutes. Table 1, which Applicants do not claim is a complete tabulation

20 of FGF receptors and species, shows certain members of the family of FGFRs and their known ligands and tissue distribution.

NAME	FGFR1		FGFR2		FGFR3		FGFR4	FGFR5
Variant	IIIb	IIIc	IIIb	IIIc	IIIb	IIIc		
Ligands	FGF-1	FGF-1	FGF-1	FGF-1	FGF-1	FGF-1	FGF-1	FGF-1
		FGF-2	FGF-2	FGF-2	FGF-2	FGF-4	FGF-2	FGF-2
		FGF-4	FGF-7	FGF-4		FGF-9		
Tissues	Brain, bone, kidney, skin, lung, heart, muscle, neuron		Brain, kidney, skin, lung, liver, glial cells		Brain, kidney, skin, lung		Lung, liver, kidney	Brain, skin, lung, testis

25

**Table 1:** FGFR variants, ligands, and tissue distribution obtained from Sigma RBI Cell Signaling and Neuroscience Catalog, 2003 (hereby incorporated by reference herein)

5

Figure 15 shows the results of a Western blot of the receptor competition experiment in Neuro-2A cells of PURE-A and either FGF-1, FGF-2, or both FGF-1 and FGF-2. Neuro-2 cells were incubated with toxin at 37°C and a concentration of 5 nM, with and without increasing concentrations of either or both FGF species. Cells were then harvested, lysed and 30 µg of protein per sample were subjected to electrophoresis in a 12% Bis-Tris polyacrylamide gel. Following transfer of the protein bands to a membrane, the membrane was probed with SMI-81 antibody that recognizes uncleaved and cleaved SNAP25. Similar experiments were performed for FGF-9, and the results were similar.

20 The competition experiment demonstrated that BoNT/A and FGF-1, FGF-2, and FGF-9 bind to the same receptor in the membrane of Neuro-2A cells. PURE-A at 5 nM without added FGF produces complete cleavage of SNAP25 to SNAP25<sub>197</sub>. It has been reported that FGF produces its physiological effects on cells at 0.1 nM, but at this concentration FGF-1 or FGF-2 do not compete with BoNT/A for receptor binding.

This result suggests that the biochemical pathway triggered by the previously characterized biological activity of FGF on cells is not sufficient to displace

30

BONT/A when bound to its receptor. Competition between FGF1 or FGF2 is first observed at an FGF concentration of 5 nM, that is, at an equimolar concentration of toxin and FGF. At a 5 nM concentration of either or both FGF1 and FGF2 cleavage of SNAP25 is reduced to about 50%. When the FGF concentration was increased to 200 nM full competition for toxin binding was achieved and all observable SNAP25 was present in its uncleaved form SNAP25<sub>207</sub>.

10

These data demonstrate that FGFR is a pathway for toxin binding and internalization. This conclusion is supported by the evidence presented here, and in light thereof, is supported by previous individual findings. For example, Schengrund et al., 29 Brain Res. Bull. 917-924 (1992) demonstrated that <sup>125</sup>I-BoNT/A binds preferentially to a protein of  $\approx$  80 kDa isolated from brain synaptosomes, and to a lesser extent to a protein of  $\approx$  116 kDa from the same extracts. These molecular weights correlate well with the bands observed in our Western blots for FGFR3. The receptor for BoNT/A is probably a glycoprotein, since binding to both the protein receptor and to the ganglioside (GT1b in the case of BoNT/A) are needed for high affinity toxin binding and internalization into neurons. In addition, the C-terminal portion of the heavy chain, that is the region involved in binding to the protein receptor, is structurally similar to FGFs.

The FGF receptors interact with complex gangliosides such as GT1b at the plasma membrane, and the activity of the receptor is modulated by these interactions (Miljan and Bremer, Science's STKE  
5 [\(<http://stke.sciencemag.org/cgi/content/full/sigtrans%3b2002/150/re15>\)](http://stke.sciencemag.org/cgi/content/full/sigtrans%3b2002/150/re15) November 26, 2002). In other systems gangliosides have been observed to modulate ligand binding, regulate receptor dimerization/polymerization, receptor activation state and subcellular localization.  
10 Neuronal cells that take up BONT/A express GT1b, and moreover, FGFR3 is expressed in spinal cord neurons during development and in motoneurons (Dono, R., 284 Am. J. Physiol. Regul. Integr. Comp. Physiol. R867-R881 (2003)). In vitro, FGFR3 ligands such as FGF2 and FGF9  
15 promote survival of spinal motoneurons in culture, and upregulate the enzyme choline acetyltransferase. Interestingly, the binding of FGF to its receptor and to BoNT/A each increase serotonin uptake by neural cells in culture (Pelliccioni et al., 17 Mol. Neurosci.  
20 303-310 (2002)).

Upon ligand binding and internalization the ligand:FGFR complexes begin their passage through endosomal compartments. The intravesicular pH drops along the endocytic pathway; this lowering of pH causes  
25 dissociation of ligand-receptor complexes and permits recycling of ligand-free receptors (Sorkin, 3 Frontiers in Bioscience d729-738 (1998)). Similarly, following binding the toxin is internalized into endosomes and after acidification of the endosome the translocation

domain (N-terminus of the heavy chain) undergoes a conformational change that facilitates delivery of the light chain to the cytoplasm. See Schiavo et al. (previously cited), Schmid et al., 394 Nature 827-830 (1993); Koriazova et al., 10 *Nat. Struct. Biol.* 13-18(2003). The FGFR ligand FGF-1 is known to be capable of crossing cellular membranes to reach the cytosol and the nucleus, and this translocation process is dependent on the maintenance of vesicle transmembrane potential (Malecki et al., 21 EMBO J. 4480-4490 (2002)).

Bafilomycin A1, a specific inhibitor of vacuolar proton pumps, blocked FGF-1 translocation completely, *id.*, and has also being described to block BoNT/A translocation to the cytoplasm from endocytic vesicles (Keller, Naunyn Schmiedeberg's Arch Pharmacol 365(Suppl 2):R26 ABS 69 (June 2002)).

This information by itself does not suggest the invention claimed herein. However, when combined with our data the cited literature demonstrate that FGFR (in the case of BONT/A, FGFR-3) functions as a selective receptor for toxin uptake and internalization.

The following examples are exemplary only and do not limit the claims, which are defined solely by their terms.

A fusion protein comprising the C terminal portion of the heavy chain of BONT and the light chain of BONT/E is tested for its ability to selectively bind and intoxicate BONT/A susceptible cells. A preparation comprising dilutions of the fusion protein is incubated with HIT-T15 insulinoma cells expressing exogenous FGFR3 in the presence of GT1b ganglioside. The ability of the fusion peptide to bind and enter the insulinoma cells is detected by detecting secretion of insulin in response to the presence of glucose, as described above. By contrast, insulin secretion is unaffected in cells not expressing FGFR3.

15

The results of this assay show that amount of insulin secreted into the culture medium is decreased in a dose-dependent manner when the fusion protein is added to the culture medium. Western blots of cell lysates will show the conversion of full length SNAP-25 to the cleaved form typical of the proteolytic activity of the BONT/E light chain protease. This assay therefore is useful in showing that the fusion peptide is able to bind and enter BONT/A susceptible cells.

25

The same fusion protein is capable of intoxicating cells of the neuromuscular junction.

30

## Example 2:

A fusion protein comprising the receptor binding portion of an FGF species capable of binding FGFR3 (including FGF1, FGF2, FGF4 and FGF9) and the translocation domain and light chain of BONT/E is tested for its ability to selectively bind and intoxicate BONT/A susceptible cells. The assay is conducted as described in Example 1 above, with similar results; the detected cleaved SNAP-25 fragments are characteristic of BONT/A intoxication.

## Example 3:

BONT/A, produced from fermentation of Clostridium botulinum is produced using standard fermentation techniques. Either or both the bulk preparation and purified, formulated versions of expressed toxin are tested for purity and activity as follows. A preparation comprising dilutions of the BONT/A preparation is incubated with HIT-T15 insulinoma cells expressing exogenous FGFR3 in the presence of GT1b ganglioside. The ability of the toxin to bind and enter the insulinoma cells is detected by detecting secretion of insulin in response to the presence of glucose, as described above. The specific activity of the preparation can be calculated from the determined protein concentration and the activity of the preparation at various doses.



These data are submitted to the U.S. Food and Drug Administration by a pharmaceutical company as part of data demonstrating how BONT/A is manufactured and tested. This information is considered by the FDA, who  
5 decides to permit the manufacture and sale of this lot of BONT/A, and subsequent lots made and tested in a similar manner, as a therapeutic pharmaceutical product based in part on this bulk and/or formulation assay data.

10

The pharmaceutical comprising the BONT/A is then offered for sale as a prescription medication.

Example 4:

15

Same as Example 3, however the polypeptide produced is the fusion neurotoxin of Example 1, produced in *E. coli*. Both bulk and/or formulation lots of the fusion neurotoxin are tested as indicated above,  
20 the data submitted to the FDA, and a decision to grant marketing approval, or continued sales of such fusion polypeptide as a therapeutic agent, is made by the FDA based at least in part on such data. The pharmaceutical company then offers the fusion  
25 neurotoxin for sale as a prescription therapeutic agent.

30

An *in vitro* assay is established using cloned FGFR3 bound to a solid support in the presence of  
5 ganglioside GT1b. The bound FGFR3 is first saturated with BONT/A heavy chain (H chain) in phosphate buffered saline (PBS), and washed free of unbound FGF. A test compound from a combinatorial library of compounds is contacted with the receptor under substantially  
10 physiological conditions (e.g., PBS), and the eluate collected. The H chain concentration in the eluate is compared to the H chain concentration of a control eluate in which H chain was not first bound to FGFR3.

15 Test compounds which are able to strongly bind FGFR3 and compete with H chain for FGFR3 binding (for example, by the method described in this section) are candidates compounds for the development of an antidote to acute botulism poisoning.

20 The examples provided herein are simply illustrations of various aspects of the invention, which is to be understood to be defined solely by the claims which follow this specification. All citations  
25 given here are hereby incorporated by reference herein.

What is claimed is:

- 5           1) A method of screening a sample for the presence  
          of an agent able to compete with BONT/A  
          neurotoxin for selective binding to cells  
          susceptible to BONT/A intoxication comprising:
  - 10           a. contacting said sample with a composition  
              comprising FGFR3 receptor and optionally GT1b  
              ganglioside, and
  - b. determining whether said molecule selectively  
              binds FGFR3,

          wherein selective binding of said molecule to

- 15           FGFR3 indicates that said molecule is able to  
          selectively compete with BONT/A neurotoxin for  
          selective binding to cells susceptible to BONT/A  
          intoxication, and wherein if said agent is BONT/A,  
          said method does not comprise an LD<sub>50</sub> assay.
- 20           2) The method of claim 1 wherein the FGFR3 is  
          expressed on the surface of a cell.
- 3) The method of claim 2 wherein said cell is
- 25           comprised in a tissue of a living animal.
- 4) The method of claim 1 wherein said contacting  
          step is performed *in vitro*.

- 5) The method of any one of claims 1-4 wherein  
said agent is BONT/A.
- 6) The method of any one of claims 1-4 wherein  
said agent is not BONT/A.
- 7) The method of claim 6 wherein said agent  
comprises a receptor binding domain of a heavy  
chain of BONT/A.
- 8) The method of claim 6 wherein said agent  
comprises a receptor binding domain of an anti-  
FGFR3 antibody.
- 9) The method of claim 6 wherein said agent  
comprises a receptor binding domain of an FGF.
- 10) The method of claim 9 wherein said FGF is  
selected from FGF 1, FGF2, FGF4 and FGF9.
- 11) A method of determining the activity of a  
preparation of BONT/A comprising the method of  
claim 1.
- 12) The method of any one of claims 1-4 or claim 1  
wherein said agent comprises an protease domain  
which cleaves a SNARE protein at a site other  
than that cleaved by BONT/A light chain.

13) The method of claim 12 wherein said protease domain comprises the active site of the light chain of a clostridial neurotoxin other than BONT/A.

5

14) The method of claim 13 wherein said protease domain comprises the active site of the light chain of BONT/E.

10

15) A method of inhibiting the activity of BONT/A in a human comprising administering to said human a first composition comprising an agent selected from the group consisting of FGFR3 and an FGF.

15

16) The method of claim 6 further comprising administering to said human a ganglioside GT1b.

20

17) A polypeptide able to compete with BONT/A for selective cell surface binding which is produced from a preparation in which the ability of said polypeptide to selectively bind to BONT/A-susceptible cells is assayed using a method comprising the steps: contacting said polypeptide with a composition comprising FGFR3 receptor and optionally GT1b ganglioside, and directly or indirectly determining whether said polypeptide selectively binds FGFR3, wherein ability to compete with BONT/A for cell surface

25

binding is indicated if said polypeptide  
selectively binds said FGFR3 receptor.

5 18) The polypeptide of claim 17 which comprises a  
cell surface binding domain of a BONT/A  
neurotoxin.

10 19) The polypeptide of claim 18 which comprises  
BONT/A.

15 20) The polypeptide of claim 18 which comprises a  
chimeric neurotoxin comprising a domain other  
than said cell surface binding domain which is  
derived from a neurotoxin other than BONT/A.

21) The polypeptide of claim 20 which comprises a  
proteolytic domain derived from BONT/E.

20 22) The polypeptide of claim 17 wherein said  
polypeptide is contacted with a receptor  
expressed on the surface of a cell.

25 23) The polypeptide of claim 22 wherein said  
ability to compete with BONT/A for cell surface  
binding is indirectly determined by detecting  
cleavage of a SNARE protein.

30 24) The polypeptide of claim 17 wherein said  
preparation is a formulated human therapeutic  
drug.

25)The polypeptide of claim 17 wherein said preparation is made from a cell lysate or culture medium.

5

26)A polypeptide comprising BONT/A which is produced from a preparation in which the ability of said polypeptide to selectively bind to BONT/A-susceptible cells is assayed using a method comprising the steps: contacting said polypeptide with a composition comprising FGFR3 receptor and optionally GT1b ganglioside, and directly or indirectly determining whether said polypeptide selectively binds FGFR3, wherein the activity of BONT/A is said preparation is directly or indirectly determined by its ability to bind FGFR3.

10

15

27)The polypeptide of claim 26 which comprises a cell surface binding domain of a BONT/A neurotoxin.

20

28)The polypeptide of claim 27 which comprises BONT/A.

25

29)The polypeptide of claim 27 which comprises a chimeric neurotoxin comprising a domain other than said cell surface binding domain which is derived from a neurotoxin other than BONT/A.

30

- 30) The polypeptide of claim 29 which comprises a proteolytic domain derived from BONT/E.
- 31) The polypeptide of claim 26 wherein said polypeptide is contacted with a receptor expressed on the surface of a cell.
- 32) The polypeptide of claim 31 wherein the activity of said BONT/A is further determined by detecting cleavage of a SNARE protein.
- 33) The polypeptide of claim 17 wherein said preparation is a formulated human therapeutic drug.
- 34) The polypeptide of claim 17 wherein said preparation is made from a cell lysate or culture medium.
- 35) The polypeptide of claim 26 in which said said ability to compete with BONT/A for cell surface binding is determined *in vitro*.
- 36) The method of claim 26 wherein said BONT/A neurotoxin comprises a native BONT/A molecule.
- 37) The method of claim 36 wherein said BONT/A molecule is isolated from a *Clostridium botulinum* strain selected from the group consisting of: CL138, 137, 129, 13, 42N, Hall



A-hyper, 667Ab, NCTC 2916, P10845, Allergan-Hall A, 62A, Kyoto-F, type A NIH, NCTC 7272, 7I03-H, and Kumgo.

5           38)A method of marketing a neurotoxin selectively  
binding to the same binding site as does BONT/A  
comprising obtaining marketing approval from a  
governmental or regional regulatory authority  
for a therapeutic neurotoxin that is first  
10           produced from a bulk preparation, wherein said  
neurotoxin is assayed for specific binding to  
neural cells using a method comprising the  
steps: contacting said neurotoxin molecule with  
a composition comprising FGFR3 receptor and  
15           GT1b ganglioside and determining whether said  
neurotoxin molecule selectively binds FGFR3,  
packaging said neurotoxin for sale in a manner  
consistent with the requirements of said  
regulatory authority, and selling said  
20           neurotoxin.

39)The method of claim 37 wherein said BONT/A  
neurotoxin comprises a native BONT/A molecule.

25           40)The method of claim 38 wherein said BONT/A  
molecule is isolated from a *Clostridium*  
*botulinum* serotype A strain selected from the  
group consisting of: CL138, 137, 129, 13, 42N,  
Hall A-hyper, 667Ab, NCTC 2916, P10845,

Allergan-Hall A, 62A, Kyoto-F, type A NIH,  
NCTC 7272, 7I03-H, and Kumgo.

- 5           41)The method of claim 38 wherein the FGFR3 is  
            expressed on the surface of a cell.
- 42)The method of claim 41 wherein said cell is  
            comprised in a tissue of a living animal.
- 10          43)The method of claim 1 wherein said contacting  
            step is performed *in vitro*.
- 44)The method of claim 38 wherein said neurotoxin  
            is not BONT/A.
- 15          45)The method of claim 44 wherein said neurotoxin  
            comprises a receptor binding domain of a heavy  
            chain of BONT/A.
- 20          46)The method of claim 44 wherein said neurotoxin  
            comprises a receptor binding domain of an anti-  
            FGFR3 antibody.
- 47)The method of claim 44 wherein said neurotoxin  
25          comprises a receptor binding domain of an FGF.
- 48)The method of claim 47 wherein said FGF is  
            selected from FGF 1, FGF2, FGF4 and FGF9.

49) The method of claim 44 wherein said neurotoxin comprises a protease domain which cleaves a SNARE protein at a site other than that cleaved by BONT/A light chain.

5

50) The method of claim 49 wherein said protease domain comprises the active site of the light chain of a clostridial neurotoxin other than BONT/A.

10

51) The method of claim 50 wherein said protease domain comprises the active site of the light chain of BONT/E.

15

52) A method of inhibiting the activity of BONT/A in a human comprising administering to said human a first composition comprising an agent selected from the group consisting of FGFR3, and anti FGFR3 antibody and an FGF.

20

53) The method of claim 6 further comprising administering to said human a ganglioside GT1b.

54) The method of claim 52 in which said agent is an anti FGFR3 antibody.

25

54) A method of screening a compound for the ability to bind to cells selectively targeted by BONT/A neurotoxin, comprising the steps contacting said compound with a composition

30

comprising FGFR3 receptor and GT1b ganglioside,  
and determining whether said compound  
selectively binds FGFR3, wherein selective  
binding of said compound to FGFR3 indicates  
5 that said molecule is able to bind to cells  
selectively targeted by BONT/A neurotoxin.

55) The method of claim 54 wherein the FGFR3 is  
expressed on the surface of a cell.

56) The method of claim 54 wherein said contacting  
step is performed *in vitro*.

57) The method of any one of claims 55 and 56  
15 wherein said molecule is BONT/A.

58) A method of rendering a cell susceptible to  
cleavage of intracellular SNARE proteins by a  
neurotoxin, comprising inducing said cell to  
20 express FGFR3.

59) The method of claim 58 wherein said induction  
step is accomplished by transfecting said cell  
with an expression vector encoding FGFR3.


60) The method of claim 59 wherein said cell is a  
pancreatic acinar cell.

61) The method of claim 59 wherein said cell is a  
30 sensory neuron.


62) The method of any one of claims 58-61 wherein  
said induction occurs in vivo.

Methods and compositions for screening ligands to  
clostridial toxin receptors. Also disclosed are  
5 methods of assaying bulk preparations of such toxin,  
and for marketing toxin so assayed.

**FIGURE 1**




**FIGURE 2A**




Control  
Electroporation  
Electroporation with PURE A






**Figure 2B**




**Figure 2C**



**Figure 3**


**BEST AVAILABLE COPY**




**Figure4:** Inhibition of insulin exocytosis caused by exogenous PURE-A on HIT-T15 cell lines. All cell lines except for control cells were previously exposed to virus containing human cDNA brain library. Cells were exposed to 1nM PURE-A. Insulin release was induced by high glucose media and was measured using an ELISA kit from Peninsula Laboratories.

**Figure 4**


Inhibition of insulin release after exposure to 1nM Pure A in  
cell lines containing the receptor for BoNT/A




**Figure 5**



**Figure 6**




**Figure 7: A.** PCR product obtained using genomic DNA from HIT-T15 C7 cell line as a template and the LIB 5' and LIB 3' primer pair. A band of 12 kDa was obtained (arrow). The band was purified and a nested PCR was performed using **WHICH???** other primer pairs. **B.** Bands of several molecular weights were obtained from the nested PCR experiment. The polynucleotide bands were purified from the gel and cloned in Topo-XL vectors for subsequent sequencing.



**Figure 8:** Western blot of 5 cell lines treated with 1 nM of Pure BoNT/A over night. Same amount of protein loaded per lane.

**A.**




**B.**




**Figure 9: A.** Time Course of SNAP25 cleavage in Neuro-2A cells treated with 1nM BoNT/A complex analyzed by Western Blot with an antibody specific for SNAP25<sub>197</sub> (different amounts of protein loaded per lane). **B.** Time Course of SNAP25 cleavage in SH-SY5Y cells treated with 1nM BoNT/A complex analyzed by Western Blot as in A.






**Figure 10:** Neuro-2a cells pre-treated with GT1b and expose to pM concentrations of toxin. 140  $\mu$ g of total protein was loaded per lane.




**Figure 11:** Overall reaction scheme to crosslink Pure A to a Putative Receptor on Neuro-2A cells (Receptor-N2A) using the cross-linking reagent sulfo-BED

**FIGURE 12**




**Figure 13**




**Figure 13:** Western blot of the toxin-receptor complex in a reducing PAGE-SDS probed with antibodies to FGFR1-4, respectively. Only antibodies to FGFR3 showed immunoreactive bands in the Neuro-2A sample.

**BEST AVAILABLE COPY**

**FIGURE 14**



**FIGURE 15**



**Figure 15:** Western blot of the competition experiment of Pure A and FGF. Toxin at 5 nM and increasing concentrations of FGF were added together to cells for 10 min at 37°C. Following electrophoresis, samples were probed in a Western blot with SMI-81 antibody, which recognizes uncleaved and cleaved SNAP25.

[illegible]

Figure 16B

	(405)	405	410	420	430	440	450	460	470	480	490	505				
BoNT/A 62A	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
BoNT/A Hall A hyper E. Johnson	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
AAC06331 BoNT/A AGN Hall A	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
I40645 BoNT/a infant bot	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
BoNT/A NTC2916	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
BoNT-A Kyoto-F	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
AAC21363 BoNT/A Kumpo	(404)	QNTETINNMFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI										
CA61124 BoNT/A 657Ab	(261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
BAA11051 BoNT/A Type ANIH	(19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16535/ BoNT/A QL138	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16534/ BoNT/A 137	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16533 BoNT/A 129	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16532 BoNT/A 13	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16531 BoNT/A 42N	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
Consensus	(405)	QNTETIN	NFTKLKNTGLPEFYKLLCVRGII	TSKTKSLD	GYNKALNDLCIKVNMWDLFFSPSEDNFTNDL	K	GEITSDTNIEAAEENISLDLIQOYYI									
	(506)	506	520	530	540	550	560	570	580	590	606					
BoNT/A 62A	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
BoNT/A Hall A hyper E. Johnson	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
AAC06331 BoNT/A AGN Hall A	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
I40645 BoNT/a infant bot	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
BoNT/A NTC2916	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
BoNT-A Kyoto-F	(505)	TFNFDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNSVNEALINSRVYTPFSSDYVKVKNA	TEAMPLG											
AAC21363 BoNT/A Kumpo	(449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
CA61124 BoNT/A 657Ab	(261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
BAA11051 BoNT/A Type ANIH	(19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16535/ BoNT/A QL138	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16534/ BoNT/A 137	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16533 BoNT/A 129	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16532 BoNT/A 13	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16531 BoNT/A 42N	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
Consensus	(506)	TF	FNDEPENISIEHLSSDIIGQL	LMPIERFPNGKKYELCKYTMFHYLRAQEFHCKSR	IALTNS	EALL	P	YTFSS	YVKVKNA	EA	MPL					
	(607)	607	620	630	640	650	660	670	680	690	707					
BoNT/A 62A	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
BoNT/A Hall A hyper E. Johnson	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
AAC06331 BoNT/A AGN Hall A	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
I40645 BoNT/a infant bot	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
BoNT/A NTC2916	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
BoNT-A Kyoto-F	(606)	WVQLVYDPTDETSEVST	DKIADITIIIPYIGPALNIGNM	LYDDFVALIPSCA	ILLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK								
AAC21363 BoNT/A Kumpo	(449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
CA61124 BoNT/A 657Ab	(261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
BAA11051 BoNT/A Type ANIH	(19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16535/ BoNT/A QL138	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16534/ BoNT/A 137	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16533 BoNT/A 129	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16532 BoNT/A 13	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16531 BoNT/A 42N	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
Consensus	(607)	W	E	LVYDPTDET	EVST	DKIADITIIIPYIGPALNIGNM	L	DFV	ALIPSG	V	LLEFIPTAIF	LPSTFALVSYIANKVLT	VQTI	NALSKRNEK		
	(708)	708	720	730	740	750	760	770	780	790	808					
BoNT/A 62A	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
BoNT/A Hall A hyper E. Johnson	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
AAC06331 BoNT/A AGN Hall A	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
I40645 BoNT/a infant bot	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
BoNT/A NTC2916	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
BoNT-A Kyoto-F	(707)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
AAC21363 BoNT/A Kumpo	(449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
CA61124 BoNT/A 657Ab	(261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
BAA11051 BoNT/A Type ANIH	(19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16535/ BoNT/A QL138	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16534/ BoNT/A 137	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16533 BoNT/A 129	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16532 BoNT/A 13	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
AAC16531 BoNT/A 42N	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----				
Consensus	(708)	DEVYKY	VTNWLAKVNTQIDLLIR	KMK	ALENQAEATRA	IINYQYNYTEEEKNN	INFNIDDLSSKLNES	IN	KAMININKFL	QC	SVSYLMNSMIF	PGVKR				
	(809)	809	820	830	840	850	860	870	880	890	909					
BoNT/A 62A	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
BoNT/A Hall A hyper E. Johnson	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
AAC06331 BoNT/A AGN Hall A	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
I40645 BoNT/a infant bot	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
BoNT/A NTC2916	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
BoNT-A Kyoto-F	(808)	LD	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD
AAC21363 BoNT/A Kumpo	(449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
CA61124 BoNT/A 657Ab	(261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
BAA11051 BoNT/A Type ANIH	(19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
AAC16535/ BoNT/A QL138	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
AAC16534/ BoNT/A 137	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
AAC16533 BoNT/A 129	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
AAC16532 BoNT/A 13	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
AAC16531 BoNT/A 42N	(1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
Consensus	(809)	L	DFDASLKID	LLKYIYDNRGTL	IGQVDR	LKIKVNTLS	DIPFQLSKYVDN	RL	LLSTPTTEY	IKNI	INTS	ILNIR	ESNHL	IDLSRYASKINIG	SKVNP	PD



FIGURE 16C

	(910)	910	920	930	940	950	960	970	980	990	1000	1010
BoNT/A 62A (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
BoNT/A Hall A hyper E. Johnson (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
AAQ06331 BoNT/A AGN Hall A (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
140645 BoNT/a infant bot (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
BoNT/A NTC2916 (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
BoNT-A, Kyoto-F (909)	-----	LDKNOIQIPNLESSK	IEVILKNAIVYNSMYENF	STSPWIRIFKYPNSI	SLNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQTQ	IEKDRVV	FKYSQMINIS	-----	-----
AAQ21363 BoNT/A Kumpo (449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
CA61124 BoNT/A 657/Ab (261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
BAA11051 BoNT/A Type A-NIH (19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16535/ BoNT/A CL138 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16534/ BoNT/A 137 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16533 BoNT/A 129 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16532 BoNT/A 13 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16531 BoNT/A 42N (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Consensus (910)	-----	LDKNOI	L NLESS	IEVILKNAIVYNSMYENF	STSPWIRIFKYP	I LNNEYTIINC	MENNSGK	VSLNYGEI	IWTLOQ	DRVV	FKYSQMINIS	-----
	(1011)	1011	1020	1030	1040	1050	1060	1070	1080	1090	1100	1111
BoNT/A 62A (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
BoNT/A Hall A hyper E. Johnson (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
AAQ06331 BoNT/A AGN Hall A (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
140645 BoNT/a infant bot (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
BoNT/A NTC2916 (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
BoNT-A, Kyoto-F (1010)	-----	YINRWIFVTITNNRIN	NSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
AAQ21363 BoNT/A Kumpo (449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
CA61124 BoNT/A 657/Ab (261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
BAA11051 BoNT/A Type A-NIH (19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16535/ BoNT/A CL138 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16534/ BoNT/A 137 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16533 BoNT/A 129 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16532 BoNT/A 13 (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16531 BoNT/A 42N (1)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Consensus (1011)	-----	YINRWIFVTITNNR	LSKIYINGRLIDQK	PISNLGNTHASNN	IMFKLDGCRD	THRYIMIKYFNL	PDKELNEKEIKDLYI	QSN	SGILKDFW	QDYLOYDK	-----	-----
	(1112)	1112	1120	1130	1140	1150	1160	1170	1180	1190	1200	1212
BoNT/A 62A (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
BoNT/A Hall A hyper E. Johnson (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ06331 BoNT/A AGN Hall A (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
140645 BoNT/a infant bot (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
BoNT/A NTC2916 (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
BoNT-A, Kyoto-F (1111)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ21363 BoNT/A Kumpo (449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
CA61124 BoNT/A 657/Ab (261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
BAA11051 BoNT/A Type A-NIH (19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16535/ BoNT/A CL138 (8)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ16534/ BoNT/A 137 (8)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ16533 BoNT/A 129 (8)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ16532 BoNT/A 13 (8)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
AAQ16531 BoNT/A 42N (8)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
Consensus (1112)	-----	YYMLNLFDPNKYVDV	NNVIGIRGYMYLKG	PRGSVVTNTIYLN	SSLYEGTKFI	LKKYASGNE	DNIVRN	NDRVYIN	VVKNKE	YRLATNAS	QAGVEKIL	SALEI
	(1213)	1213	1220	1230	1240	1250	1260	1270	1280	1297		
BoNT/A 62A (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
BoNT/A Hall A hyper E. Johnson (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
AAQ06331 BoNT/A AGN Hall A (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
140645 BoNT/a infant bot (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
BoNT/A NTC2916 (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
BoNT-A, Kyoto-F (1212)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS
AAQ21363 BoNT/A Kumpo (449)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
CA61124 BoNT/A 657/Ab (261)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
BAA11051 BoNT/A Type A-NIH (19)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16535/ BoNT/A CL138 (78)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16534/ BoNT/A 137 (78)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16533 BoNT/A 129 (78)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16532 BoNT/A 13 (78)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AAQ16531 BoNT/A 42N (78)	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Consensus (1213)	-----	PDVGNLSQVVMKSK	DDQGITNKCKMNL	DDNNGNDIGF	IGPHF	DNIAKL	VASNNY	NRQIER	SSRTL	GCSWEF	IPVDDG	WGESS

BEST AVAILABLE COPY